



## **Blood transfusion practices at Al-Thawra Teaching Hospital, Sana'a-Yemen.**

Ahmed kaid Salem, MD

Department of Medicine, Faculty of Medicine and Health sciences. Department of Medicine, Al-Thawra Hospital

### **Abstract**

*Objectives: To evaluate the blood transfusion practices, the quality and quantity and the use rate of blood and blood components among patients at Al-Thawra Teaching Hospital.*

*Methods: This a descriptive study that has been conducted for the purpose of studying the blood transfusions fate and practices for all crossmatched units in the blood bank at Al-Thawra hospital during one month, from 31/12/2002-31/01/2003. Data were collected from blood bank and from different departments.*

*Results: 960 units were crossmatched during the period of the study. The study showed that crossmatched units for male patients (56%) were higher than for female (44%). The recipient male/female ratio was 1.27: 1. The mean age of recipients was 31.6 years± SD. The commonest type of blood was blood group O (48%). 87.5% of crossmatched blood was done for stored whole blood in spite of availability of other components. Medical indications represented the highest percentage (40%). 26% of all crossmatched blood units were returned to the bank. The majority of collected blood was by family donation system. It was obvious that there was over-ordering and irrational use of the blood from the high percentage of unused crossmatched blood and the use of stored whole blood.*

*Conclusion: Blood transfusion is necessary for saving the life of the patient who is in need, but blood should be only given when necessary, to minimize its complications and to cut off extra cost for cross matching of unneeded blood.*

*The collection of blood from voluntary non-remunerated blood donors from low- risk populations should be encouraged. Modernization of blood banks with automated separator of the blood components is mandatory.*

**Key words:** Blood, blood bank, blood components, donors, recipients, Yemen

### **Introduction:**

Blood transfusion is an old method of restoring person believed to be dying by passing blood by another into his vein. The earliest attempt of blood treatment was an effort to save the life of the dying Pope Innocent VIII in 1492 having him drink the blood of 3 healthy boys. (1, 2)

Although, the association of blood with life and vitality was recognized by primitive man, the transfusion of blood was not possible until Harvey described the circulation in 1628. (3)

The era of blood transfusion started with the discovery of

the ABO blood group system by Landsteiner in 1931. In addition, his discovery of Rhesus factor in 1940 provided an explanation for previously mysterious hemolytic reactions in occasional blood recipients. (4)

This technique of red cells antibody detection, lead to recognition of many new blood group antigen system and introduction of reliable cross-matching of patient and donor blood.

Nowadays, transfusion medicine has become a fundamental part of medical practice in all medical disciplines and different specialties. The major function of

**Corresponding author: To Dr. Ahmed Kaid Salem, Hematology/Oncology senior Consultant at Al-Thawra hospital. PO Box 20552 Sana'a. Republic of Yemen. Tel. 733760019. Email: ahmdkaid@yahoo.com**

the hospital transfusion service is promptly providing appropriate quantity of safe blood and blood components to patients in critical needs.

Requirement of blood and blood component is increasing day by day but the number of donors doesn't increase correspondingly, hence, our hospitals face shortage of blood periodically. Furthermore, when a unit of blood is cross-matched for a particular patient, it is temporarily removed from the blood inventory and is unavailable for use by another patient, thereby leading to a shortage of blood in the blood bank. Also, cross-matching is not an inexpensive procedure. It requires costly reagents, equipment, and manpower. So avoidance of unnecessary cross-matching is essential to cut cost. (5) Although modern technology has made blood transfusion a generally safe procedure; a definite risk is involved so that transfusion should be given thereby only when the benefit to the patient exceeds the inherent danger to the procedure. While advances have been dramatic in most industrialized countries, blood transfusion in the developing countries has tended to stagnate at post II World War level, with chronic shortage, lack of component therapy and unsolved safety problems.(1)

In Yemen the blood transfusion services still in the primitive stage, in spite of that they provide hospitals with a large quantity of requested blood.

At Al-Thawra Teaching Hospital the Blood Transfusion Unit is approximately daily, receiving 80 blood requests, collecting 70 Units of blood and delivering 70 Units of blood.

Despite the lack of previous study in this field, in Yemen, this study seems to be very important because the need for blood and blood component is increasing day by day, due to life modernization. Also there is a paucity of donors, because of lack of awareness of this vital national problem leading to shortage of blood and its components. Also lack of modern equipments in blood banks which depend mainly on family donor system all over the country makes the blood insufficient and unsafe.

### Objectives of the study:

-General objective: to highlight the blood transfusion practices in Al-Thawra Teaching Hospital.

### -Specific objectives:

- 1.To determine the relative distribution of cross-matched blood in the different departments of the hospital.
- 2.To evaluate the use rate of blood and blood components among patients who need blood.
- 3.To determine the crossmatched to transfused ratio (C/T) and the resulted extra-amount of unneeded cross-matched blood.
- 4.To ensure that the components required are made available in both quantity and quality.

### Material and Methods:

The study is a descriptive study for whole cross-matched units of blood, done in the blood bank at Al-Thawra Hospital during the period 31.12.2002 to 31.01.2003.

Data collection started at blood bank then completed in the different departments of the hospital through the patients and their files. The following information was filled into a structured questionnaire from the request paper including:

- a) Name, age, and sex of the patients.
- b)Department from which the blood was ordered.
- c)Blood group and Rh of cross-matched blood units.
- d) Availability of requested blood.
- e)Methods of blood dispersion from blood bank (relative donors, free or by guarantee).
- f) Pre-transfusion tests.
- g)Indications of blood transfusion. The indications were classified into: medical, surgical, traumatic and gynecological & obstetrical regardless of the department.
- h) Site of blood transfusion: The hospital was categorized into 5 departments: medical, surgical, gynecological & obstetrical, pediatric and general emergency & emergency ICU.
- i)Fate of the cross-matched units, whether if they are used or returned to the blood bank. The time between delivering and returning the blood was recorded.


Data analysis:

Variables were recorded in one chart. Variables were inserted and analyzed by (S.P.S.S. program). (7) The obtained results are discussed & represented. Finally according to the results we state the conclusions, which were the base for our recommendations.

**Results:**

**I. Demographic data:**

960 cross-matched units of blood were studied. Information regarding age, sex, blood group and Rh was available for the recipients. 538 (56%) units were cross-matched for male patients. 422 (44%) units were cross-matched for female patients. The male/ female ratio was 1.27:1. Figure (1)



The mean age of patients for whom the cross-matching tests were done was 31.6 ±18.3 "median 30" years.


Table (1): Shows the age groups of the recipient patients and as appear in this table most patients were under the age of 50 years (65%). (87.5%) of patients were under the age of 70 years.

Table (1): Frequency of cross-matched units among age groups

Age group	Frequency	Percentage
1-<10	115	12%
10-<20	123	12.8%
20-<30	123	12.8%
30-<40	102	10.6%
40-<50	166	17.3%
50-<60	97	10.1%
60-<70	114	11.9%
70-<80	120	12.5%

Most of the blood group cross-matched during the period of the study were as shown in figure (2) O+ve, 457 (48%), and the least patients were B-ve, one patient only.

Figure (2): Blood groups distribution.



**II. Blood bank service:**

During the period of study all blood groups were available except only 4 units were not available at the blood bank (three units were AB-ve and one unit was AB+ve) and these were requested from other hospitals.

The necessary pre-transfusion tests such as: VDRL, for syphilis, HbsAg for hepatitis B, HCV antibodies for hepatitis C and HIV antibodies for AIDS were done for all units and written on all bags before cross-matching.


**III. Component of blood available in blood bank:**

Three types of blood & blood components were available:

- Whole blood (stored and fresh)
- Frozen plasma(stored & fresh)
- Packed RBCs

Most blood requests were for whole blood 842 (87.5%) and the smallest number for stored plasma16 (1.7%).


Figure (3): Types of blood components of cross-matched units.



#### IV. Departments:

960 units were cross-matched for patients in different departments of the hospital. As shown in figure (4) most of the units were cross-matched for patients in surgical ward 551 (58%) units, for different indications, and the least cross-matched units, 82 (8.5%) were for gynecological and obstetrical department.

Figure (4): Distributions of cross-matched units by departments.



#### V. Indications for blood transfusion

The indications for transfusion were classified into 4 categories regardless the department from which the patient came. During the period of the study we found that the highest frequency for crossmatched units were for medical indications 387 (40%) units, and the smallest number of crossmatched units were for gynecological and obstetrical indications 76 (8%) units, see figure (5).





Figure (5): Indications for blood transfusion.

The most common medical indication, as shown in figure (6) was anemia 212 (55%) units.

Figure (6) Medical indications for blood transfusion



Surgical indications for transfusion rank the second. The highest number of crossmatched units were for patients whom they will undergo cardiac surgery 147 (41%) units, and the least number was for pediatric surgery 10(3%) units, see figure (7).




Figure (7): Surgical indications for blood transfusion.

Traumatic indications for blood transfusion ranked the third, 142 (15%) units, as shown in figure (5). Most crossmatched units for traumatic indications were for patients admitted to the hospital after road traffic accident and gunshot injury 65 (46%) units for each, and the smallest number was for patients with stab wound injury only two units (1.4%). See figure (8)




Figure (8): Traumatic indications for blood transfusion

Gynecological and obstetrical indications for blood transfusion, were the least common indication and ranked the fourth, 76 (8%) units, see figure (5). The most common crossmatched units were for gynecological indications 49 (64.5%) units, e.g. dysfunctional uterine bleeding & hysterectomy. The rest were for obstetrical indications, 27 (35.5%) units.

#### VI. Returned blood:

During the period of the study we found that from whole crossmatched blood 247 (26%) units were returned to blood bank and 222 (90%) units of the returned blood were returned after leaving the blood bank immediately, (transfusion was cancelled) 25 (10%) units were returned after some time of delivering them from the blood bank. The median time of delay between delivering the blood and returning it to the blood bank was approximately 3 hours.

The high percentage of returned blood (38%) was from the surgical ward, whereas the lowest percentage of returned blood was from the medical and gynecological & obstetrical wards (13%) for each.

#### VII. Blood dispersion from the blood bank:


The blood was given to the patients by three methods:

1.580 (61%) units were given for the patients by guarantee until the patient gives donors to the bank. Most of the units given by guarantee were for patients in surgical ward (55%) and the least number for the pediatric ward (6%).

2.221(23%) units were given to the patients who previously brought donors to the blood bank before

transfusion. Most of these units were for patients in the surgical ward (92%) and the smallest number was for patients in the medical ward (0.5%).

3.156 (16%) units were given free to the patients, without guarantee or donors. The majority of the units were for patients in the medical ward (40%) and the smallest percentage (3%) was for patients in gynecological and obstetrical ward. See figure (9).



#### Discussion:

This is the first study as we know that was designed to follow the fate of blood transfusion at Al-Thawra Teaching Hospital and may be in Yemen as a whole. In our daily practice we may observe some over prescribing of blood and irrational use of the blood. Therefore this study was conducted to look more closely to this issue.

In our study, most of the crossmatched units were for male patients (56%), in contrast to study that was done in North England by Wells A.W. et al, nearly half (49.3%) of the crossmatched units were for female patients. (8) This increase in male patients who need blood, in our country, is due to male persons are the active part of the population and they may face trauma and/or stress which may necessitate blood transfusion.

The median age of patients for whom crossmatching tests were done was 30 years. This result is similar to that reported from South Africa (9). While in the study that done by Wells the median age of patients was 67 years. (8) In other study done in Sweden in 1993, the median age was 71 years and only 10% of patients were under the age

of 50 years and (38%) of the crossmatched were for patients under the age of 70 years. (10)

In our study, however, (65.5%) of patients were under age of 50 years and (87.5%) were under age of 70 years. Thus, there is a big difference between our country and Western countries regarding the age of patients who are in need for blood. This difference may be due to young population pyramid of our country in which young people form most of the society. In contrast to old population pyramid of Western countries where the percentage of old people is high. In addition, the increase in the incidence of all forms of trauma in our active young people may reflect this difference.

In this study, the most common blood groups for which crossmatching tests were done were blood group O and A, found in (54.5%) and (34%) respectively. Whereas the blood group B was infrequent. The majority of the patients were Rh+ (90%) and a minority was Rh- (10%). These results approximately are similar to that recorded in Saudi Arabia and in Caucasoid. (11, 12, 13)

From this study it was clear that the most common indication was the medical indication (40%) followed by surgical (37%), traumatic (15%), and the least common indication for blood was the gynecological and obstetrical indication (8%).

These results are in accordance with North England study which found (51.6%) for medical indications followed by surgical (40.7%) and small percentage were for gynecological and obstetrical indications. However, other study in Sweden showed that the highest percentage of indications was for surgical followed by medical in (47%) and (29%) correspondingly. (10)

In North England study the traumatic indications were only (3%) of all indications (8), compared to 15% in our study. This, indicate that trauma is more frequent in our country than in England.

#### **Blood components:**

The preparation of blood components allows a single blood donation to provide treatment for two or three

patients and also avoids the transfusion of blood elements of the whole blood that patient may not require. (14)

Whole blood should only be used when increased oxygen carriage and massive circulatory volume expansion are required, which is a rare situation. Also stored whole blood should be assumed not to possess any worthwhile haemostatic function. (14)

Our study reveals that despite of presence of other blood components to a considerable degree in the blood bank of the hospital, most of the blood requests, during the study period were for stored whole blood (87.5%) for different indications. The percentage of requested other components of blood like packed RBCs, and plasma was very small. This result is in contradiction with blood transfusion practices worldwide where the whole blood is requested in small number of patients who need blood volume replacement. (15, 16, 17)

As it was noticed in patients with anemia that actually need the oxygen-carrying capacity of red cells and not the plasma for volume replacement, most of the blood requests were for whole stored blood (87.3%) and only small percentage (16.7%) for red blood. This practice, represent a major irrational use of blood.

We think that the cause of not ordering blood components frequently, despite its availability in the blood bank, is that the separation of the blood components is usually done manually which is rudimentary and not effective at all. Thus, the separation of the blood components should be done by automatic machine and stored in suitable conditions.

Although, blood was screened for only four known blood transmitted diseases (HIV, HBV, HCV and syphilis), we can not exclude other blood transfused diseases for which no screening was done e.g. malaria, HTLV-1 and others not known and no tests are available to detect them.

Also the viremia, but antibody negative "window period" remains a major problem globally. But the automated genomic amplification techniques such as Polymerase Chain Reaction (PCR) to individual donations may be able

to resolve this problem with little regard for high costs. However, near-absolute safety of blood cannot be ensured through technical/ laboratory solutions alone without regard for all the other vital and complementary elements which are essential to quality and safety. Such as collection of blood only from selected voluntary non-remunerated donors and judicious clinical use of blood. (18)

The actual study revealed that the over-ordering of the blood at our hospital is obviously clear from the high percentage of returned blood (26%) which was not needed and so lead to shortage of blood in the blood bank. When a unit of blood is crossmatched for a particular patient it is temporarily removed from the bank capacity and is made unavailable for using by another patient, who may need it at that time. The blood bank keeps the crossmatched blood reserved for "48" hours. Most of the returned blood (90%) was returned to the bank after, median time of "3" hours, making this blood unsafe due to hemolysis and contamination.

Also the cross-matching tests cost the patient 1000 rials for each unit, so avoidance of this unnecessary corssmatching is essential to cut this cost.

The highest percentage of returned blood was from the surgical wards (38%) that reflect the over-ordering of the unnecessary blood for surgical patients.

#### **Blood dispersion from the bank:**

In our study, the donation of blood was mainly depending on family donation system. This means that the relatives are donating for a particular patient. This type of blood donation is abandoned worldwide and is replaced by voluntary non-remunerated donors. In Taiwan, since 1974 all donations depend on this system. (19) Consequently, there are major problems if there is no organized blood transfusion service and no coordinated plan for a modern blood transfusion service. Since the unavailability of blood in a life-threatening condition is obviously a catastrophic situation for the patient and his relatives to make the blood available immediately. And

since the organization of a volunteer donor program requires sophisticated and long-term arrangements, the problem is approached in the only possible way, by making the patient, family and friends responsible for providing donors.

The majority of cross-matched units were given for patients by guarantee for the bank (61%), which obligates the patient to bring 2 donors for each unit. That means they must replace double amount, making this blood unsafe for recipients, because the obligated family will arrange this even from high risk group of donors. 23% of blood was given to patients who have relatives' donors before. There are several reasons why this "family donor system" is unsatisfactory, unsafe and dangerous:

- It may not be possible for patient's family to find suitable donors and the family may feel obligated to donate even if they know that they have some health problem which prohibits blood donation.

This may also encourage "blood selling practice" from whom Hb level is low and as well as the risk of certain transmissible disease is higher. Although it is well known that certain category in Yemen can sell the blood, due to poverty and khat or drugs addiction, we were unable to investigate this issue in our study.

It is not possible, therefore, to utilize family donation system effectively for creating a blood bank, nor is it possible to utilize such blood for effective production of platelets and other blood components.

- Finally, since this blood is dedicated to a particular patient, it may often be transfused to that patient even if not ultimately needed.

Only 16% of blood was given free of charge, which the correct way for blood transfusion.

#### **Conclusion:**

The stored whole blood was the commonest blood component transfused in spite of availability of some blood components, like packed RBCs and plasma. This type of transfusion leads to exhaustion of the blood storage and expose the patient to many complications that

can be avoided if selected component needed by patient is given.

Over ordering of the blood which appear from the high percentage of unused crossmatched units, leads to shortage and exhaustion of the bank and expose the blood to hemolysis which becomes unsuitable for transfusion. Also make high cost for patients and the bank.

The most common indication for blood transfusion was medical indication, regardless of the ordering ward. Also the study showed high percentage of traumatic indications mostly in male.

The absence of suitable policy for blood collection and dispersion in the blood bank and the dependency mostly on the "family donor system" as main source for blood, which obligate the patient to bring donors to the bank, cause extra cost for patient and his/her family. Also the family donors may feel obligated to donate even if they know that they have some health conditions which prohibits donation of blood. Consequently "Family donor system" is unsafe and dangerous blood could be collected.

#### Recommendations:

Conducting similar research, for a longer period, in different hospitals.

Spreading of our findings among health workers, with stress on over ordering of the blood and the type of the blood use.

Health education about rational use of the blood. Providing the blood bank with modern equipment for preparation and preservation of blood components.

Organization of a volunteer donor program and not to depend fundamentally on family donor system as a main source for the blood.

The establishment of nationally coordinated blood transfusion services with quality system in hospitals over the country.

#### Acknowledgment:

The authors would like to thank the hematology group of students of the 15th batch, who were the cornerstone of the study. Also, our best thanks to the staff of the hospital

particularly staff of the blood bank that, helped us during our study. Also, we would like to thank our entire colleague who helped us in one way or in another.

#### References:

1. Britten A.F., Ala F.A, El-Nageh M (1994). Blood transfusion: A Basic Text Book, WHO Regional office for the Eastern Mediterranean.
2. Jayaprakash P.A., Al-Azzawi S. (2001): Maximum surgical blood order schedule. A safe, effective and economic solution to preoperative over-ordering of blood; Oman Med J; 18(2):11-16.
3. Beulter E. (2001): Preservation and clinical use of blood and blood components. Hematology Ed by William J Williams, 6th edition. MCGRAW-HILL Publishing Company; PP 1879-1892.
4. Mollison PL: (1983). Red cell antigens and antibodies and their interactions. In Blood transfusion in clinical medicine. 7th ed. By Blackwell scientific publications, Oxford London; PP 191: 268.
5. Behrman R.E, Vaughan V.C & Nelson W.E (1987): Transfusion. Nelson Textbook of Pediatrics. 13th ed. PP 1059-1061. W.B. SAUNDERS Company.
6. Gunson H., Martlew V.J (1996): Blood transfusion. Weatherall, Ledingham & Warrell (Eds): Oxford Text Book of Medicine. 3rd Ed, (3) PP: 3687-3695. Oxford university press, Oxford, New York and Tokyo.
7. Statistical Package for Social Sciences-Personal Computers (SPSS-PC). Version III. Illinois: Chicago SPSS Co; 1996.
8. Wells A.W, Mounter P.J, Chapman C.E, et all (2002): Where does blood go? Prospective observational study of red cell transfusion on north England, BMJ; 325(7368):803-804.
9. Patricia K (2003): Transfusion medicine: Education in the developing world. Africa Sanguine, (6); 3-5.
10. Tynell E, Norde R, Shanwell A, et all (2001): Long-term survival in transfusion recipients in Sweden, 1993; transfusion, 41(2):252-55.

11. Guyton C (1996): Blood cells, immunity and blood clotting. In Hall J.E eds. Textbook of medical physiology, 19th ed, by W.B. Saunders Company, PP425-430.
12. Bashwari L.A, Al-Mulhim A, Ahmed M.S, and Ahmed MA: (2001) Frequency of ABO blood groups in the eastern region of Saudi Arabia. Saudi Med J, (22): 1008-1012.
13. Sukkar MY, El-Munshid HA, Ardawl MSM. In: Blood. In Concise Human Physiology. Oxford, UK Blackwell Scientific Publications; 1993.
14. Menitove J.E. (1996): Blood transfusion. In, Bennet, Plum, and Eds: CECIL Textbook of Medicine, 20th Ed. By W.B. Saunders Company. PP893-897.
15. The Clinical use of blood-Handbook. WHO-Geneva, 2001.
16. Vincent J.L, Baron J.F, Reinhart K, Gattinoni L, and et all: (2002): Anemia and blood transfusion in critically ill patients. JAMA; 288: 1499-1507.
17. Hoffbrand AV, Pettit JE and Moss PAH. (2001). Blood transfusion. In: Essential Haematology. 4th ed. Blackwell Science Publications. PP 307-318.
18. Heyns AD (2000): The risk of transmitting HIV and other diseases with blood transfusion in South Africa. Transfusion Today, 42; 4-9.
19. Burnouf T, Cuthberston B, Lin L, Liu DT et all (2000): Experience in implementing collection of plasma for fractionation in Taiwan. Transfusion today, 42; 9-13.